

Predicate Calculus

Simon Foster & Jim Woodcock

University of York

PROF

11th December 2021

Overview

Introduction

Reasoning with Quantifiers

De Morgan's Laws

Quantifier Laws

Summary

Outline

Introduction

Reasoning with Quantifiers

De Morgan's Laws

Quantifier Laws

Summary

Introduction

- ▶ Predicate quantifiers: extent that proposition is true over range of elements.
- ▶ Predicate = proposition with abstracted subject: $P(x)$.
- ▶ **Universal Quantification** Predicate true for all the values over domain.
- ▶ $\forall x \bullet P(x)$: states that $P(x)$ is true for all values of x in this domain.
- ▶ **Existential Quantification** Predicate true for some value of x in the domain.
- ▶ $\exists x \bullet P(x)$: states that $P(x)$ is true for at least one value in the domain.

Syntax

- ▶ In Z, quantifiers share a similar syntax: $\exists x : s \mid P \bullet Q$.
- ▶ \exists is the **quantifier**, x is the **bound variable**, and s is the **range** of x .
- ▶ P is the (optional) constraint and Q is the predicate.
- ▶ Two syntactic equivalences explain the constraint:

$\exists x : s \mid P \bullet Q$ is a shorthand for $\exists x : s \bullet P \wedge Q$

$\forall x : s \mid P \bullet Q$ is a shorthand for $\forall x : s \bullet P \Rightarrow Q$

- ▶ $\exists x : s \mid P \bullet Q$ read as “there exists an x in s satisfying P , such that Q ” holds.
- ▶ $\forall x : s \mid P \bullet Q$ read as “for all x in s satisfying P , Q holds”.

Bound Variables

- ▶ Each quantifier introduces a **bound variable**.
- ▶ Scoped, like local variables in a block-structured programming language.
- ▶ Consider the quantified predicate “ $\forall x : s \mid P \bullet Q$ ”.
- ▶ Bound variable **scope**: exactly the constraint P and predicate Q : $\forall x : s \mid \underbrace{P \bullet Q}_{\substack{\text{scope} \\ \text{of } x}}$.
- ▶ Quantifiers take the widest possible scope, binding very loosely: $\forall x : s \bullet P \wedge Q$.
- ▶ This **means** “ $\forall x : s \bullet (P \wedge Q)$ ”. **Not** “ $(\forall x : s \bullet P) \wedge Q$ ”.
- ▶ Variables not yet bound are **free**. Example: $x > 3$ has one free variable: x .
- ▶ Truth-tables are **useless** for quantifiers, in general.
- ▶ Because bound variables can range over sets that are too large to tabulate.

Substitution

- ▶ Suppose P is a predicate containing free variable x .
- ▶ “ $\forall x : s \bullet P$ ”: asserts P is true for every x .
- ▶ Specific theorems are obtained by **substituting** a term t for bound variable x .
- ▶ $P[t/x]$ denotes P with term t substituted for x . Read as “ P with t for x ”.

Some authors use replacement notation: $P[x \setminus t]$, $P[x := t]$.

Remember your English grammar: Substitute new for old. Replace old by new.

- ▶ Substitution also defined on terms themselves: $u[t/x]$.
- ▶ In Isabelle, we write $P(x)$ to show that x is free for substitution in P .
- ▶ Substitution is done by supplying an actual parameter: $P(t)$.

Substitution

- ▶ Bound variables must sometimes be renamed before substitution.
- ▶ Consider the theorem $\forall x : \mathbb{Z} \bullet \exists y : \mathbb{Z} \bullet x \neq y$.
- ▶ True for all numbers, so why not specialise it? Is it true for y : $\exists y : \mathbb{Z} \bullet y \neq y$?
- ▶ **Contradiction!** — What's gone wrong? Free variable **capture**.
- ▶ **Free** variable y enters scope of $\exists y : \mathbb{Z} \bullet \dots$ and becomes a **bound** variable.
- ▶ Change the bound variable before substitution to avoid capture:

$$(\exists y : \mathbb{Z} \bullet x \neq y)[y/x] = (\exists z : \mathbb{Z} \bullet x \neq z)[y/x] = \exists z : \mathbb{Z} \bullet y \neq z$$

Outline

Introduction

Reasoning with Quantifiers

De Morgan's Laws

Quantifier Laws

Summary

Reasoning with Quantifiers

- ▶ Variable capture example suggests: the name of a bound variable doesn't matter.
- ▶ Predicates that differ up to renaming of bound variables are equivalent.

$$\exists y : \mathbb{Z} \bullet x \neq y = \exists z : \mathbb{Z} \bullet x \neq z$$

- ▶ What about other equivalences in the predicate calculus?
- ▶ Here's a simple question about two quantified predicates:

Are these two predicates equivalent?

$$\exists x : s \bullet P(x) \Rightarrow Q(x) \quad ?= \quad (\exists x : s \bullet P(x)) \Rightarrow (\exists x : s \bullet Q(x))$$

- ▶ Are they equivalent? Does one imply the other? Are they at all related?
- ▶ We need identities to reason about quantifiers to check the answer.
- ▶ Before that, we need rules for reasoning about quantifiers.

Reasoning with Universal Quantification

- ▶ Suppose $P(x)$ is true for every possible value of x in s : then $\forall x : s \bullet P(x)$.
- ▶ Otherwise, $\forall x : s \bullet P(x)$ is false.
- ▶ This gives us an argument.
- ▶ **Generalisation:** $x \in s, P(x) \vdash \forall x : s \bullet P(x)$
- ▶ This is an argument with two premises and one conclusion.
- ▶ This argument is **valid** only if its premises are true for every x .
- ▶ A new kind of restriction: the variable x must be **arbitrary**.
- ▶ **Valid:** $x \in \mathbb{Z}, x + 1 > x \vdash \forall x : \mathbb{Z} \bullet x + 1 > x$
- ▶ **Fallacy:** $x \in \mathbb{Z}, x = 5 \vdash \forall x : \mathbb{Z} \bullet x = 5$

Reasoning with Universal Quantification

- ▶ If $\forall x : s \bullet P(x)$ is true, then for any term t in s , $P(t)$ will be true.
- ▶ This gives us an argument.
- ▶ **Specialisation:** $\forall x : s \bullet P(x), t \in s \vdash P(t)$
- ▶ **Example:** $\forall x : \mathbb{Z} \bullet x^2 \geq 0, y - 1 \in \mathbb{Z} \vdash (y - 1)^2 \geq 0$
- ▶ This is another argument with two premises and one conclusion.

Reasoning with Existential Quantification

- ▶ Suppose $P(x)$ is true for at least one element of s : $\exists x : s \bullet P(x)$.
- ▶ Otherwise, $\exists x : s \bullet P(x)$ is false.
- ▶ This gives us an argument: $t \in s, P(t) \vdash \exists x : s \bullet P(x)$
- ▶ **Existential introduction**. This hides information by giving a name to a witness.
- ▶ **Existential elimination**: prove a predicate R from the predicate $\exists x : s \bullet P(x)$.
- ▶ Show that R follows from an **arbitrary** element of s satisfying P .

$$\exists x : s \bullet P(x), (x \in s, P(x) \vdash R) \vdash R$$

- ▶ Similar to generalisation (universal introduction).
- ▶ Note the sub-argument $x \in s, P(x) \vdash R$ as a premise to the overall argument.
- ▶ How are existential elimination and universal introduction linked?
- ▶ De Morgan's laws for quantifiers.

Outline

Introduction

Reasoning with Quantifiers

De Morgan's Laws

Quantifier Laws

Summary

De Morgan's Laws for Quantifiers: Motivation

- Can the predicate $\neg \forall x : \mathbb{N} \bullet x > 3$ be simplified? **Semi-formal argument:**

$$\begin{aligned} & \neg \forall x : \mathbb{N} \bullet x > 3 \\ \Leftrightarrow & \quad \{ \text{universal quantification as conjunction} \} \\ & \neg (0 > 3 \wedge 1 > 3 \wedge 2 > 3 \wedge 3 > 3 \wedge \dots) \\ \Leftrightarrow & \quad \{ \text{by De Morgan's Law} \} \\ & \neg (0 > 3) \vee \neg (1 > 3) \vee \neg (2 > 3) \vee \neg (3 > 3) \vee \dots \\ \Leftrightarrow & \quad \{ \text{existential quantification as disjunction} \} \\ & \exists x : \mathbb{N} \bullet \neg (x > 3) \end{aligned}$$

- Motivates De Morgan's Laws for generalised conjunction and disjunction:

$$\neg (\forall x : s \bullet P(x)) \Leftrightarrow \exists x : s \bullet \neg P(x) \quad \text{and} \quad \neg (\exists x : s \bullet P(x)) \Leftrightarrow \forall x : s \bullet \neg P(x)$$

Example: A Classic Logic Puzzle

- ▶ Suppose that if you don't love someone, then you hate them.
- ▶ Is there no one who loves everybody? Draw conclusions if it's true.
- ▶ Let's calculate. Let *Person* be the set of all people:

$$\begin{aligned} & \neg \exists x : \textit{Person} \bullet \forall y : \textit{Person} \bullet x \textit{ loves } y \\ \Leftrightarrow & \quad \{ \text{by De Morgan} \} \\ & \forall x : \textit{Person} \bullet \neg \forall y : \textit{Person} \bullet x \textit{ loves } y \\ \Leftrightarrow & \quad \{ \text{by De Morgan} \} \\ & \forall x : \textit{Person} \bullet \exists y : \textit{Person} \bullet \neg (x \textit{ loves } y) \\ \Leftrightarrow & \quad \{ \text{by definition} \} \\ & \forall x : \textit{Person} \bullet \exists y : \textit{Person} \bullet x \textit{ hates } y \end{aligned}$$

- ▶ So, our sentence means everybody hates someone.

Example: De Morgan for Quantifiers

- ▶ For every pair of integers x and y , there's a z , such that $x + z = y$.
- ▶ Formalised as: $\forall x : \mathbb{Z} \bullet \forall y : \mathbb{Z} \bullet \exists z : \mathbb{Z} \bullet x + z = y$ (\mathbb{Z} is the set of **integers**)
- ▶ Or more simply as: $\forall x : \mathbb{Z}; y : \mathbb{Z} \bullet \exists z : \mathbb{Z} \bullet x + z = y$.
- ▶ Or even more simply as: $\forall x, y : \mathbb{Z} \bullet \exists z : \mathbb{Z} \bullet x + z = y$.
- ▶ Predicate is **false** for all positive integers in \mathbb{N} : $\forall x, y : \mathbb{N} \bullet \exists z : \mathbb{N} \bullet x + z = y$.
- ▶ To see this, take $x = 2$ and $y = 1$, then we must prove that $\exists z : \mathbb{N} \bullet 2 + z = 1$.
- ▶ Which is patently **false**: there is no such z . $\exists z : \mathbb{N} \bullet z = -1$.
- ▶ How do we prove it's a contradiction?

Let's Calculate

It's false. Negate the predicate and show that the result is true.

$$\begin{aligned} & \neg \forall x, y : \mathbb{N} \bullet \exists z : \mathbb{N} \bullet x + z = y \\ \Leftrightarrow & \quad \{ \text{by De Morgan} \} \\ & \exists x, y : \mathbb{N} \bullet \neg \exists z : \mathbb{N} \bullet x + z = y \\ \Leftrightarrow & \quad \{ \text{by De Morgan} \} \\ & \exists x, y : \mathbb{N} \bullet \forall z : \mathbb{N} \bullet \neg (x + z = y) \\ \Leftrightarrow & \quad \{ \text{by definition} \} \\ & \exists x, y : \mathbb{N} \bullet \forall z : \mathbb{N} \bullet x + z \neq y \end{aligned}$$

We can prove our result more easily in this form.

Finishing it off

- So how do we prove $\exists x, y : \mathbb{N} \bullet \forall z : \mathbb{N} \bullet x + z \neq y$?

- Use existential introduction!

- Find values for x and y that make the following true

$$\forall z : \mathbb{N} \bullet x + z \neq y$$

- If we select 1 for x and 0 for y , we have

$$\forall z : \mathbb{N} \bullet 1 + z \neq 0$$

which is a property of every natural number: $\text{succ}(z) \neq 0$.

- So, given this property, we've proved that

$$\neg \forall x, y : \mathbb{N} \bullet \exists z : \mathbb{N} \bullet x + z = y$$

More on De Morgan

- What happens to the range of the bound variable? Consider the following:

$$\begin{aligned} & \neg \forall x : \mathbb{N} \bullet x^2 + x - 2 > 0 \\ \Leftrightarrow & \quad \{ \text{since } \forall x : s \bullet P \text{ is the same as } \forall x : X \bullet x \in s \Rightarrow P \} \\ & \neg \forall x : \mathbb{Z} \bullet x \in \mathbb{N} \Rightarrow x^2 + x - 2 > 0 \\ \Leftrightarrow & \quad \{ \text{by De Morgan} \} \\ & \exists x : \mathbb{Z} \bullet \neg (x \in \mathbb{N} \Rightarrow x^2 + x - 2 > 0) \\ \Leftrightarrow & \quad \{ \text{by propositional calculus} \} \\ & \exists x : \mathbb{Z} \bullet x \in \mathbb{N} \wedge x^2 + x - 2 \leq 0 \\ \Leftrightarrow & \quad \{ \text{since } \exists x : s \bullet P \text{ is the same as } \exists x : X \bullet x \in s \wedge P \} \\ & \exists x : \mathbb{N} \bullet x^2 + x - 2 \leq 0 \end{aligned}$$

- Range of the quantification remains unchanged.

Outline

Introduction

Reasoning with Quantifiers

De Morgan's Laws

Quantifier Laws

Summary

Proving and Disproving Universal Quantifications

- ▶ Proving $\forall x : s \bullet P(x)$ can be quite demanding.
- ▶ We need an argument that proves P whatever the value of x .
- ▶ Not enough to give some examples of values of x that satisfy P .
- ▶ Disproving $\forall x : s \bullet P(x)$ may be much easier.
- ▶ By De Morgan, it's the same as proving $\exists x : s \bullet \neg P(x)$.
- ▶ We need only a single x for which P is false.
- ▶ Value for x provides a counterexample to $\forall x : s \bullet P(x)$.

Example: Disproving a Universal Quantification

- ▶ Show the following statement is false

$$\forall x : \mathbb{Q} \mid x > 0 \bullet (x^2 - 3x + 2 \geq 0)$$

\mathbb{Q} is the rational numbers (those expressible as the ratio of two integers a/b).

- ▶ A single counterexample suffices.
- ▶ An appropriate value is $\frac{3}{2} > 0$.
- ▶ We have that

$$\left(\frac{3}{2}\right)^2 - 3 * \left(\frac{3}{2}\right) + 2 = -\frac{1}{4} < 0$$

Moving Quantifier Scopes

- ▶ Suppose our predicate is a disjunct or conjunct within a quantifier.
- ▶ Suppose none of its variables are bound by the quantifier.
- ▶ Then the predicate can be pulled out of the quantifier scope.
- ▶ Let N stand for a predicate in which x doesn't occur free.
- ▶ We have the following equivalences

$$\forall x : s \bullet P(x) \wedge N \Leftrightarrow (\forall x : s \bullet P(x)) \wedge N$$

$$\forall x : s \bullet P(x) \vee N \Leftrightarrow (\forall x : s \bullet P(x)) \vee N$$

$$\exists x : s \bullet P(x) \wedge N \Leftrightarrow (\exists x : s \bullet P(x)) \wedge N$$

$$\exists x : s \bullet P(x) \vee N \Leftrightarrow (\exists x : s \bullet P(x)) \vee N$$

Moving Quantifiers Around

- ▶ Suppose we have a predicate that is a universally quantified conjunction.
- ▶ Suppose the bound variable occurs in both parts of the conjunction.

$$\forall x : s \bullet P(x) \wedge Q(x)$$

- ▶ Can we move the quantifier inwards?
- ▶ Distributivity laws for the quantifiers

$$(\forall x : s \bullet P(x) \wedge Q(x)) \Leftrightarrow (\forall x : s \bullet P(x)) \wedge (\forall x : s \bullet Q(x))$$

$$(\exists x : s \bullet P(x) \vee Q(x)) \Leftrightarrow (\exists x : s \bullet P(x)) \vee (\exists x : s \bullet Q(x))$$

- ▶ Maybe they're really associativity laws?

Example: Distributivity

- **WARNING:** Universal quantification doesn't distribute over disjunction:

$$\neg \left(\left(\forall x : s \bullet P(x) \vee Q(x) \right) \Leftrightarrow \left(\forall x : s \bullet P(x) \right) \vee \left(\forall x : s \bullet Q(x) \right) \right)$$

- Counterexample to distribution:

Every number is either even or odd:

$$\forall n : \mathbb{Z} \bullet (n \bmod 2 = 0) \vee (n \bmod 2 = 1)$$

But it's false that either every number is even, or every number is odd:

$$\left(\forall n : \mathbb{Z} \bullet (n \bmod 2 = 0) \right) \vee \left(\forall n : \mathbb{Z} \bullet (n \bmod 2 = 1) \right)$$

Example: Distributivity

- **WARNING:** Existential quantification doesn't distribute over conjunction:

$$\neg \left((\exists x : s \bullet P(x) \wedge Q(x)) \Leftrightarrow (\exists x : s \bullet P(x)) \wedge (\exists x : s \bullet Q(x)) \right)$$

- Counterexample to distribution:

There's a number that's even and there's a number that's odd:

$$(\exists n : \mathbb{Z} \bullet (n \bmod 2 = 0)) \wedge (\exists n : \mathbb{Z} \bullet (n \bmod 2 = 1))$$

But there isn't a number that's both even and odd:

$$\exists n : \mathbb{Z} \bullet (n \bmod 2 = 0) \wedge (n \bmod 2 = 1)$$

Semi-distribution

- ▶ Universal quantification doesn't distribute over disjunction as an equivalence.
- ▶ But it does as an implication (**strengthening**):

$$(\forall x : s \bullet P(x)) \vee (\forall x : s \bullet Q(x)) \Rightarrow (\forall x : s \bullet P(x) \vee Q(x))$$

- ▶ Existential quantification doesn't distribute over conjunction as equivalence.
- ▶ But it does as an implication (**weakening**)

$$(\exists x : s \bullet P(x) \wedge Q(x)) \Rightarrow (\exists x : s \bullet P(x)) \wedge (\exists x : s \bullet Q(x))$$

Identities and Semi-identities

1. $(\forall x : s \bullet P(x)) \Rightarrow P(c)$ $c \in s$
2. $P(c) \Rightarrow (\exists x : s \bullet P(x))$ $c \in s$
3. $(\neg \exists x : s \bullet P(x)) \Leftrightarrow (\forall x : s \bullet \neg P(x))$
4. $(\forall x : s \bullet P(x)) \Rightarrow (\exists x : s \bullet P(x))$
5. $(\neg \forall x : s \bullet P(x)) \Leftrightarrow (\exists x : s \bullet \neg P(x))$
6. $(\forall x : s \bullet P(x) \wedge N) \Leftrightarrow (\forall x : s \bullet P(x)) \wedge N$
7. $(\forall x : s \bullet P(x) \vee N) \Leftrightarrow (\forall x : s \bullet P(x)) \vee N$
8. $(\forall x : s \bullet P(x)) \wedge (\forall x : s \bullet Q(x)) \Leftrightarrow (\forall x : s \bullet P(x) \wedge Q(x))$
9. $(\forall x : s \bullet P(x)) \vee (\forall x : s \bullet Q(x)) \Rightarrow (\forall x : s \bullet P(x) \vee Q(x))$
10. $(\exists x : s \bullet P(x) \wedge N) \Leftrightarrow (\exists x : s \bullet P(x)) \wedge N$
11. $(\exists x : s \bullet P(x) \vee N) \Leftrightarrow (\exists x : s \bullet P(x)) \vee N$
12. $(\exists x : s \bullet P(x) \wedge Q(x)) \Rightarrow (\exists x : s \bullet P(x)) \wedge (\exists x : s \bullet Q(x))$
13. $(\exists x : s \bullet P(x)) \vee (\exists x : s \bullet Q(x)) \Leftrightarrow (\exists x : s \bullet P(x) \vee Q(x))$

Example: Multiple Quantifiers

- ▶ Order of universal and existential quantifiers is significant.
- ▶ “No matter what value x in s , a value y in t can be found such that. . .”

$$\forall x : s \bullet \exists y : t \bullet \dots$$

- ▶ Value of y may depend on value of x . **Example:** “there’s a larger one”

$$\forall x : \mathbb{N} \bullet \exists y : \mathbb{N} \bullet y > x$$

(1)

- ▶ “There’s a value y in t so that no matter what value x in s is chosen, . . .”

$$\exists y : t \bullet \forall x : s \bullet \dots$$

- ▶ Since y is bound first, y must be specified independently of x .
- ▶ “There’s a number that’s greater than every other number”:

$$\exists y : \mathbb{N} \bullet \forall x : \mathbb{N} \bullet y > x$$

(2)

- ▶ Which is true and which is false?

Outline

Introduction

Reasoning with Quantifiers

De Morgan's Laws

Quantifier Laws

Summary

Summary

- ▶ Previously: **Propositional logic**: truth-valued propositions.
- ▶ Truth functional connectives:
 - conjunction, disjunction, negation, implication, equivalence.
- ▶ **First-order predicate logic** adds predicates and quantification.
- ▶ **Predicates**: parametrised propositions with abstracted subjects.
- ▶ **Quantifiers**: universal and existential: for all \dots , exists at least one \dots
- ▶ Quantifiers apply to variables in parametrised predicates:
 - bound, free, substitution.
- ▶ Rich set of algebraic laws.
- ▶ **Next lecture**: Extending the natural deduction system to quantifiers.